

WEEKLY NEW BRUNSWICK INFLUENZA REPORT

Reporting period: April 6 to April 12 2014 (week 15)

Summary:

In New Brunswick, low influenza activity in week 15, all positive detections were influenza B.

New Brunswick:

- There have been 12 positive influenza detections during week 15, all were influenza B.
- The ILI consultation rate decreased slightly compared to previous week and was within the expected levels for this time of year.
- One new influenza outbreak was reported in a nursing home in region 3.

Canada:

- In week 15, influenza activity in Canada was sustained by continued circulation of influenza B. Ontario reported a marked increase in influenza activity and some western provinces have also reported increases in detections of influenza B. Influenza activity remains within the expected levels for this time of year.
- While the influenza A (H1N1)pdm09 virus has mostly affected adults 20-64 years of age this season, influenza B is having a greater impact on adults 65 years of age and older, as well as young persons 5 to 19 years of age.
- 722 laboratory detections of influenza were reported in week 15. The percentage of laboratory tests positive for influenza was 15.3%.
- The national ILI consultation rate was 30.7 consultations per 1,000 patients' visits, which is within the expected range for week 15.
- Nine new influenza outbreaks were reported: all in long-term care facilities. Also, 2 influenza outbreaks were reported in other settings and 1 outbreak of ILI was reported in a school.

International:

- Human infection with Avian Influenza: As of April 16 2014, a total of 421 laboratory-confirmed cases of human infection with an avian influenza A (H7N9) virus were reported in China (as well as in Taiwan, Hong Kong and Malaysia) including 145 deaths. The majority of cases have presented with severe acute illness, rapidly progressing to severe pneumonia. Most human cases have reported a history of exposure to poultry or live bird markets. There is currently no evidence of sustained human-to-human transmission of H7N9.
- MERS-CoV: Since April 2012, 243 laboratory-confirmed cases have been reported from Saudi Arabia, Qatar, Jordan, United Arab Emirates, Kuwait, United Kingdom, Oman, France, Germany, Tunisia, Italy and Malaysia. Among the 243 cases, 93 were fatal. Onset of illness was between April 2012 and April 2014.
- Novel influenza A viruses:
 - Since summer 2013, the United States reported 21 new cases of human infection with variant influenza A viruses (19 H3N2v and 2 H1N1v) from Illinois, Indiana, Ohio, Michigan, Arkansas and Iowa. No human-to-human transmission has been identified. All have reported close contact with swine.
 - China reported 3 human cases of avian-origin influenza A(H10N8) in recent months (with exposure to live poultry markets) with 2 deaths. While human infection with other H10 subtypes, notably H10N7, has been previously reported, these are the first reports of H10N8 infection in humans.

1) Influenza Laboratory Data¹

- Low influenza activity, all positive detections were influenza B.
- 12 influenza detections were reported during this current reporting period.
- Since the beginning of the season, 1392 positive influenza detections were reported, 440 influenza A (H1N1)pdm09, 1 influenza A (H3), 884 influenza A (unsubtyped) and 67 influenza B.

Surveillance specimens are submitted by recruited New Brunswick Sentinel Practitioner Influenza Network (NB SPIN) practitioners, which are comprised of 8 sites in Emergency Rooms, 3 sites in Family Practice, 2 sites in First Nations communities, 1 site in a Nursing Home, 3 sites in Universities and 8 sites in Community Health Centers. Diagnostic specimens are submitted by physicians in the community/hospital setting. Influenza laboratory data is comprised of results from surveillance and diagnostic specimens. All laboratory specimens are tested using a real-time PCR assay, which is a rapid detection method designed for detection of all known variants of influenza A and B. All laboratory-confirmed cases are reported for the week when laboratory confirmation was received.

<u>Graph 1</u>: Number and percent of positive influenza specimens in New Brunswick by week, up to April 12 2014 (data source: G. Dumont Lab results)




Table 1: Positive influenza test results by Health Region, in New Brunswick up to April 12 2014 (data source: G. Dumont lab results)


Region	Reporting period: Apr./06/2014–Apr./12/2014						Cumulative: (2013/2014 season) Aug./25/2013 –Apr./12/2014					Cumulative: (2012/2013 season) Aug./26/2012 – Aug./24/2013					
		А				В		АВВ				АВВ					
	Activity level ²	A(H1)	A(H3)	(H1N1) pdm09	unsubt yped		Total	A(H1)	A(H3)	(H1N1) pdm09	unsubt yped		Total	Non- (H1N1) pdm09	(H1N1) pdm09		Total
Region 1	Sporadic	0	0	0	0	3	3	0	1	205	442	15	663	527	13	18	558
Region 2	No activity	0	0	0	0	0	0	0	0	86	219	1	306	211	3	8	222
Region 3	Localized	0	0	0	0	1	1	0	0	41	80	3	124	85	9	1	95
Region 4	Sporadic	0	0	0	0	6	6	0	0	52	61	31	144	168	5	3	176
Region 5	No activity	0	0	0	0	0	0	0	0	10	22	5	37	20	1	7	28
Region 6	Sporadic	0	0	0	0	1	1	0	0	42	49	9	100	252	5	50	307
Region 7	Sporadic	0	0	0	0	1	1	0	0	4	11	3	18	89	2	11	102
Tota	Total NB		0	0	0	12	12	0	1	440	884	67	1392	1352	38	98	1488

² Influenza activity level definition is available on the PHAC FluWatch website: http://www.phac-aspc.gc.ca/fluwatch/13-14/def13-14-eng.php

2) ILI Consultation Rates³

- During week 15, the ILI consultation rate was 10.9 consultations per 1,000 patient visits, and was within the expected levels for this time of year.
- During week 15, the sentinel response rate was 38% for both the FluWatch sentinel physicians and the NB SPIN practitioners.

Graph 2: ILI Consultation Rates in New Brunswick, by report week, season 2013/14 compared to previous seasons*



^{*} The mean rate was based on data from the 1996/97 to 2012/2013 seasons and excludes the Pandemic season (2009-2010).

3) ILI and Laboratory-Confirmed Outbreak Data

<u>Table 3</u>: ILI activity/outbreaks in New Brunswick nursing homes and schools for the reporting week, current and previous seasons.


	А	Reporting period: pr./06/2014–Apr./12/201	Cumulative # of	Cumulative # of outbreaks		
	Lab-confirmed outbreaks in Nursing Homes*	Schools reporting ILI outbreaks**	Lab-confirmed outbreaks in Other Settings*	outbreaks season 2013-2014	season 2012-2013	
Region 1	0 out of 13	0 out of 74	0	3	15	
Region 2	0 out of 15	0 out of 81	0	2	38	
Region 3	1 out of 14	0 out of 95	0	4	20	
Region 4	0 out of 6	0 out of 22	0	1	2	
Region 5	0 out of 2	0 out of 18	0	0	6	
Region 6	0 out of 9	0 out of 35	0	3	23	
Region 7	0 out of 4	0 out of 27	0	2	10	
Total NB	1 out of 63	0 out of 352	0	15	114	

^{*}Two or more ILI cases within a seven day period, including at least one laboratory-confirmed case of influenza. Outbreaks are reported in the week when laboratory confirmation is received.

^{**}Schools reporting greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or Public Health Authority) which is likely due to ILI.

³ A total of 34 practitioner sites (19 FluWatch sentinel physicians and 15 NB SPIN sites) are recruited this season to report the number of ILI patients and total patient consultations one day during a reporting week.

<u>Graph 3</u>: Number of Influenza Outbreaks in Nursing Homes¹ and ILI Outbreaks in Schools² reported to Public Health in New Brunswick, by report week, season 2013/14.



¹ The National FluWatch definition of an outbreak in a nursing home is stated as two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case.

National Flu Watch Program - Additional information on influenza activity in Canada and around the world is available on the Public Health Agency of Canada's website at: http://www.phac-aspc.gc.ca/fluwatch/

Other Links:

World-http://www.who.int/influenza/surveillance monitoring/updates/latest update GIP surveillance/en/index.html

Europe: http://www.euroflu.org/cgi-files/bulletin_v2.cgi and

http://www.ecdc.europa.eu/en/healthtopics/seasonal influenza/epidemiological data/Pages/Weekly Influenza Surveillance Overview.aspx

PAHO: http://new.paho.org/hq/index.php?option=com_content&task=blogcategory&id=805&Itemid=569]

Australia: http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm]

New Zealand: [http://www.surv.esr.cri.nz/virology/influenza weekly update.php

Argentina: http://www.msal.gov.ar/ South Africa: http://www.nicd.ac.za/ US: www.cdc.gov/flu/weekly/

> Prepared by the Communicable Disease Control Unit Office of the Chief Medical Officer of Health, Tel: (506) 444-3044

² The National FluWatch definition of an ILI outbreak in a school is stated as absenteeism greater than 10% (or absenteeism that is higher (e.g.>5-10%) than expected level as determined by school or Public Health Authority) which is likely due to ILI.