

# WEEKLY NEW BRUNSWICK INFLUENZA REPORT

Reporting period: April 1 to April 7 2018 (week 14)

### Summary:

## In New Brunswick, most influenza indicators remained elevated in week 14

#### New Brunswick:

- There have been 122 positive influenza cases in week 14. To date this season, 2193 cases have been reported, 240 had influenza A (H3), 17 had influenza A (H1N1)pdm09, 876 had influenza A (unsubtyped, most likely the H3 strain), 1048 had influenza B and 12 had both influenza A and B simultaneously.
- Both Influenza A(H3N2) and Influenza B are co-circulating this season.
- There have been 10 new influenza associated hospitalizations during week 14. So far this season, 542 influenza associated hospitalizations have been reported with 29 deaths.
- The ILI consultation rate was 17.1 consultations per 1,000 patients visits in week 14. The ILI rate was within the expected levels for this time of year.
- Two new influenza outbreaks were reported in week 14. So far this season, 36 outbreaks were reported in total: 5 in hospitals, 19 in nursing homes, 4 in other settings (special care homes and adult residential facilities) and 8 ILI outbreaks in schools.

#### Canada:

- Influenza activity in Canada continued to decrease, but many parts of the country are still reporting localized activity. All indicators of influenza activity decreased from the previous week.
- Detections of influenza B are similar to those of influenza A.
- To date this season, the majority of lab confirmations, hospitalizations and deaths with influenza have been among adults aged 65 years and older.

#### International:

### Seasonal influenza:

• Influenza activity decreased in most of the countries in the temperate zone of the northern hemisphere, with exception of Eastern Europe where activity continued to increase. In the temperate zone of the southern hemisphere, influenza activity remained at inter-seasonal levels. Worldwide, influenza A and influenza B accounted for a similar proportion of influenza detections.

### Effectiveness of 2017-2018 influenza vaccine for influenza A(H3N2):

- In September 2017, the <u>WHO indicated</u> that the effectiveness of vaccines containing A/Hong Kong/4801/2014 was suboptimal in the 2017 southern hemisphere season, in regions where A(H3N2) viruses predominated. If influenza A(H3N2) viruses predominate in the 2017/2018 northern hemisphere season, it is likely that vaccine effectiveness would be suboptimal for the A(H3N2) virus; however, the vaccine should provide good protection for influenza A(H1N1)pdm09 and influenza B virus infection.
- In the context of a potentially reduced influenza VE for the 2017-18 season, the Association of Medical Microbiology and Infectious Disease (AMMI Canada) has posted an updated <u>quidance on the use of antiviral medication</u>.

## **Emerging Respiratory Viruses:**

- MERS CoV:
  - o WHO: http://www.who.int/csr/disease/coronavirus infections/en/
  - o CDC: <a href="http://www.cdc.gov/coronavirus/mers/">http://www.cdc.gov/coronavirus/mers/</a>
- Avian Influenza:
  - o WHO: www.who.int/csr/disease/avian\_influenza/en/index.html


### 1) Influenza Laboratory Data<sup>1</sup>

- Most influenza indicators remained elevated in week 14.
- One-hundred-twenty-two influenza cases were reported during week 14.

<sup>&</sup>lt;sup>1</sup> Surveillance specimens are submitted by recruited New Brunswick Sentinel Practitioner Influenza Network (NB SPIN) practitioners, which are comprised of sites in Emergency Rooms, in Family Practice, in First Nations communities, in Nursing Home, in Universities and in Community Health Centers. Diagnostic specimens are submitted by physicians in the community/hospital setting. Influenza laboratory data is comprised of results from surveillance and diagnostic specimens. All laboratory specimens are tested using a real-time PCR assay, which is a rapid detection method designed for detection of all known variants of influenza A and B. All laboratory-confirmed cases are reported for the week when laboratory confirmation was received.

Since the beginning of the season, 2193 influenza cases were reported, 240 with influenza A (H3), 17 with influenza A(H1N1)pdm09, 876 with influenza A (unsubtyped)<sup>2</sup>, 1048 with influenza B and 12 with co-infection of influenza A & B simultaneously.


<u>Graph 1</u>: Number and percent of positive influenza specimens<sup>3</sup> in New Brunswick by week, up to April 7 2018 (data source: G. Dumont Lab results)



<sup>&</sup>lt;sup>2</sup> The influenza A (unsubtyped) detections are most likely of the predominant strain (H3).

<sup>&</sup>lt;sup>3</sup> Total number of positive influenza tests is higher than number of cases since some individuals had co-infection of A & B simultaneously.

Figure 2: Influenza/ILI activity levels<sup>4</sup> by Health Zones, in New Brunswick, for reporting week, season 2017/2018.



<sup>4</sup> No activity is defined as no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported. Sporadic activity is defined as sporadically occurring ILI and lab confirmed influenza detection(s) with no outbreaks detected within the influenza surveillance region.

<sup>&</sup>lt;u>Localized activity</u> is defined as evidence of increased ILI with lab confirmed influenza detection(s) and outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in less than 50% of the influenza surveillance region.

<sup>&</sup>lt;u>Widespread activity</u> is defined as evidence of increased ILI with lab confirmed influenza detection(s) and outbreaks in schools, hospitals, residential institutions and/or other types of facilities occurring in greater than or equal to 50% of the influenza surveillance region.


<u>Table 1</u>: Positive influenza cases by Health Region, in New Brunswick for reporting week, cumulative current and previous seasons. (data source: G. Dumont lab results up to April 7 2018)

	Reporting period:						Cumulative: (2017/2018 season)						Cumulative: (2016/20167 season)					
	April/01/2018–April/07/2018						Aug./27/2017 –Apr./07/2018						(2016/20167 Season) Aug./28/2016 –Aug./26/2017					
Zone	А				В	A & B co- infection	А				A & B co- infection	А			В	A & B co- infection		
	A(H3)	(H1N1) pdm09	Unsubty ped/ Other	A Total	Total	Total	A(H3)	(H1N1) pdm09	Unsubty ped/ Other	A Total	Total	Total	(H3)	(H1N1) pdm09	Unsubty ped/ Other	A Total	Total	Total
Zone 1	0	4	20	24	28	2	83	. 7	407	497	653	8	75	0	505	579	89	1
Zone 2	2	0	4	6	15	0	26	1	102	129	125	1	21	1	77	99	8	0
Zone 3	3	1	5	9	19	0	51	2	167	220	94	3	25	0	117	142	23	0
Zone 4	0	0	0	0	0	0	19	2	53	74	84	0	18	0	31	50	5	1
Zone 5	0	0	1	1	0	0	8	0	8	16	6	0	2	0	3	5	6	0
Zone 6	1	0	1	2	4	0	38	3	74	115	45	0	27	0	62	89	11	0
Zone 7	1	0	3	4	8	0	15	2	65	82	41	0	21	0	52	73	16	0
Total NB	7	5	34	46	74	2	240	17	876	1133	1048	12	189	1	845	1037	158	2

### ILI Consultation Rates<sup>5</sup>

- During week 14, the ILI consultation rate was 17.1 consultations per 1,000 patients visits. The ILI rate was within the expected levels for this time of year.
- During week 14, the sentinel response rate was 26%, for both the FluWatch sentinel physicians and the NB SPIN practitioners.

Graph 2: ILI Consultation Rates in New Brunswick, by report week, season 2017/18 compared to previous seasons\*



<sup>\*</sup> The mean rate was based on data from the 1996/97 to 2016/2017 seasons and excludes the Pandemic season (2009/10).

### 3) ILI and Laboratory-Confirmed Outbreak Data

<u>Table 2</u>: ILI activity/outbreaks in New Brunswick nursing homes and schools for the reporting week, current and previous seasons.

	April	Reporting period: /01/2018–April/07/2018	Cumulative # of outbreaks	Cumulative # of outbreaks		
	Lab-confirmed outbreaks in Nursing homes <sup>6</sup>	ILI school outbreaks <sup>7</sup>	Lab-confirmed outbreaks in Other settings <sup>4</sup>	season 2017-2018	season 2016-2017	
Zone 1	0 out of 13	0 out of 74	1	6	3	
Zone 2	0 out of 16	0 out of 81	0	9	5	
Zone 3	1 out of 14	0 out of 95	0	14	14	
Zone 4	0 out of 6	0 out of 22	0	1	0	
Zone 5	0 out of 2	0 out of 18	0	0	1	
Zone 6	0 out of 9	0 out of 35	0	3	0	
Zone 7	0 out of 4	0 out of 27	0	3	2	
Total NB	1 out of 64	0 out of 352	1	36	25	


<sup>&</sup>lt;sup>5</sup> A total of 28 practitioner sites (16 FluWatch sentinel physicians and 12 NB SPIN sites) are recruited this season to report the number of ILI patients and total patient consultations one day during a reporting week.

5

<sup>&</sup>lt;sup>6</sup> Two or more ILI cases within a seven day period, including at least one laboratory-confirmed case of influenza. Outbreaks are reported in the week when laboratory confirmation is received.


Schools reporting greater than 10% absenteeism which is likely due to ILI.

<u>Graph 3</u>: Number of Influenza Outbreaks (nursing homes, hospitals, other) and ILI Outbreaks (schools) reported to Public Health in New Brunswick, by report week, season 2017/18.



## 4) Influenza associated Hospitalization<sup>8</sup> and Death<sup>9</sup> Surveillance<sup>10</sup>

<u>Graph 4</u>: Influenza associated Hospitalizations and ICU admissions in New Brunswick, by week of hospitalization for current and past season.\*



<sup>\*</sup>Those who had been hospitalized 15 days or more prior to laboratory confirmation date were excluded from the graph

\*\* Twenty-nine deaths have been reported so far in season 2017-2018.

 $<sup>^{8}</sup>$  Hospitalizations (including ICU admissions) are influenza associated; they may or may not be due to influenza.

<sup>&</sup>lt;sup>9</sup> Deaths are influenza associated; influenza may not be the direct cause of death.

<sup>&</sup>lt;sup>10</sup> In early January 2014, the Office of the Chief Medical Officer of Health implemented a new provincial surveillance system in collaboration with the Regional Health Authorities to monitor influenza-associated hospitalizations, intensive care unit admissions and deaths. A standardized Enhanced Surveillance Form is used to collect data on hospitalizations.

National Flu Watch Program - Additional information on influenza activity in Canada and around the world is available on the Public Health Agency of Canada's website at: <a href="http://www.phac-aspc.gc.ca/fluwatch/">http://www.phac-aspc.gc.ca/fluwatch/</a>

### Other Links:

World-http://www.who.int/influenza/surveillance\_monitoring/updates/latest\_update\_GIP\_surveillance/en/index.html

Europe: http://www.ecdc.europa.eu/en/healthtopics/seasonal\_influenza/epidemiological\_data/Pages/Weekly\_Influenza\_Surveillance\_Overview.aspx

PAHO: <a href="http://new.paho.org/hq/index.php?option=com\_content&task=blogcategory&id=805&Itemid=569">http://new.paho.org/hq/index.php?option=com\_content&task=blogcategory&id=805&Itemid=569</a>]

Australia: <a href="http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm">http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-surveil-ozflu-flucurr.htm</a>]

New Zealand: [http://www.surv.esr.cri.nz/virology/influenza\_weekly\_update.php

Argentina: <a href="http://www.msal.gov.ar/">http://www.msal.gov.ar/</a>
South Africa: <a href="http://www.nicd.ac.za/">http://www.nicd.ac.za/</a>
US: <a href="http://www.nicd.ac.za/">www.cdc.gov/flu/weekly/</a>

Prepared by the Communicable Disease Control Unit Office of the Chief Medical Officer of Health, Tel: (506) 444-3044